



Contribution ID: 378

Type: **Talk in the parallel session**

## Chiral vortical effect for free fermions on anti-de Sitter space

*Monday, 5 July 2021 17:30 (20 minutes)*

According to the axial vortical effect, an axial current  $J_A^\mu$  is produced in a fluid undergoing a macroscopic vortical motion, which is equal to the local kinematic vorticity  $\omega^\mu$  multiplied by the axial vortical conductivity  $\sigma_A^\omega$ . We probe the curvature corrections to  $\sigma_A^\omega$  by computing the thermal expectation value of  $J_A^\mu$  with respect to a rigidly-rotating quantum state at finite temperature. The calculation is computed in the real time formalism using a novel KMS relation which includes the effect of rotation, being based on an exact expression for the fermion vacuum two-point function (the analysis is restricted to subcritical rotations when no speed of light surface forms, such that the rotating and stationary vacua are identical). Our results confirm the Minkowski expression for  $\sigma_A^\omega$ , revealing a novel contribution proportional to the Ricci scalar. At vanishing mass, the conservation of  $J_A^\mu$  implies a non-vanishing flux through the adS boundary, while at non-vanishing mass, the flux of  $J_A^\mu$  is completely converted into a volumetric density of pseudoscalar condensate  $-i\bar{\psi}\gamma^5\psi$ .

**Primary author:** AMBRUS, Victor E. (Goethe University)

**Co-author:** Prof. WINSTANLEY, Elizabeth (University of Sheffield)

**Presenter:** AMBRUS, Victor E. (Goethe University)

**Session Classification:** Quantum Fields

**Track Classification:** Early Universe: Quantum Fields