



Contribution ID: 551

Type: Talk in the parallel session

Binary neutron star formation and the origin of GW170817

Friday 9 July 2021 07:45 (25 minutes)

The first neutron star-neutron star merger (NS-NS: GW170817) was detected in gravitational waves by LIGO/Virgo in a galaxy in which the majority of star formation was taking place a long time ago (~ 11 Gyr). Only some extreme evolutionary models (with small NS natal kicks and high common envelope efficiency) can generate NS-NS merger rates in old host galaxies consistent with the LIGO/Virgo estimate ($> 100 \text{ Gpc}^{-3} \text{yr}^{-1}$). However, we show that these models generate rates exceeding empirical Galactic NS-NS merger rates based on the large population of Milky Way NS-NS binaries. Typically, evolutionary models produce NS-NS merger rates that are consistent with the Milky Way empirical rates ($\sim 10\text{--}200 \text{ Myr}^{-1}$). However, these models generate local cosmic NS-NS merger rate in old host galaxies ($\sim 1\text{--}70 \text{ Gpc}^{-3} \text{yr}^{-1}$) below the LIGO/Virgo estimate. The reason behind this tension is the predicted delay time distribution between star formation and NS-NS mergers that favors short delays. Evolutionary models produce a generic steep power-law NS-NS delay time distribution. This limits NS-NS merger rates in old host galaxies. However, we show that such distribution is consistent with observations of Galactic NS-NS binaries; 50% of which show very long merger times (much longer than Hubble time). Once model distributions are convolved with continuous prolonged star formation in the Galactic disk, then $\sim 20\text{--}70\%$ of the predicted NS-NS population has very long current Galactic merger times (> 30 Gyr). Although NS-NS binaries are formed predominantly with short delay times, many of short delay time systems merge, while long delay time systems survive and contribute to the current Galactic NS-NS population. This study highlights the tension between the current evolutionary predictions and the observation of the first NS-NS merger in an old host galaxy. It is crucial to understand that models need to explain not only the LIGO/Virgo rate estimate, but also the merger site.

Authors: BELCZYNSKI, Krzysztof; Prof. BULIK, Tomasz; OLEJAK, Aleksandra (Nicolaus Copernicus Astronomical Center Polish Academy of Sciences); CHRUSLINSKA, Martyna; SINGH, N.; POL, N.; ZDUNIK, L.; O' SHAUGHNESSY, R.; MCLAUGHLIN, M.; LORIMER, D.; KOROBKIN, O.; VAN DEN HEUVEL, E.P.J.; DAVIES, M.B.; HOLZ, D.

Presenter: OLEJAK, Aleksandra (Nicolaus Copernicus Astronomical Center Polish Academy of Sciences)

Session Classification: Explosive Events Associated with Compact-Object Binary Mergers

Track Classification: Binaries: Explosive events associated with compact-object binary mergers