

Testing Λ CDM with eBOSS


Ryan Keeley

Marcel Grossman 2021


Testing the Concordance Model

- Λ CDM + GR *
- Λ - test via low-redshift distances *
- CDM - test via small scale structure
- GR - test via growth rate measurements *
- Inflation - CMB, LSS
- Testing FLRW (homogeneity + isotropy)

Accelerated Expansion of the Universe



H0 Tension





- Inferences from the CMB predict $H(z=0) = 67.36 \pm 0.54$ km/s/Mpc
- Measuring H0 directly gives 74.03 ± 1.42 km/s/Mpc
- Difference is now at $4.4\text{-}\sigma$.
- No obvious systematics
- Potentially a challenge for Λ CDM

Case for low- z solutions

- The physics at the CMB is non-trivial
- Thus, it is natural to expect adding new physics between the CMB and today would solve the H_0 tension
- Such solutions inherit the high-redshift successes of Λ CDM
- CMB does not measure H_0 only predicts it
- The CMB only constrains H_0 via the constraint on $\theta_s = r_s(z_*)/D_A(z_*)$
- Geometric degeneracy \rightarrow there exists $w(z)$ such that the CMB is well fit and predicts any* value of H_0

Guardrails

- BAO measure both $D_A(z)$ and $H(z)$
- 5 tracers in 7 redshift bins $z \sim 0.1$ to 2.3
- The curves are predictions from just the Planck data (not fits to the eBOSS data)
- SN measure luminosity distances $D_L(z)$
- 1048 SN from $z \sim 0.01$ to 2.3
- Both datasets are unanchored and thus cannot tell which value of H_0 is correct
- Can only constrain a mutual scale $H_0 r_d$
- They can constrain the possible expansion histories that map between $z=0$ and $z=z^*$



Throwing everything at the wall

- The strategy is to throw whatever extensions to LCDM we can think of at the datasets and see if anything sticks
- Curved CPL - $w(z) = w_0 + w_a \frac{z}{1+z}$
- Chebyshev polynomials $w(z) = - \sum_1^4 c_i T_i(x)$, $x = \log(1+z)/\log(1+z_*)$
- GP regression
- If these very broad cases cannot resolve the H0 tension, then we must conclude low-z physics as a whole cannot solve the H0 tension

GP

- Gaussian process - a distribution of functions characterized by a covariance function

$$\langle \gamma(s_1)\gamma(s_2) \rangle = \sigma_f^2 e^{-(s_1-s_2)^2/(2\ell^2)}$$

- hyperparameters σ_f and ℓ control heights and lengths of the random fluctuations respectively.

GP

- An instance or a sample of a GP, $\gamma(z)$, is a hyperfunction that randomly varies around the “mean function”, H_{mf} , $\Rightarrow H_i(z) = \gamma_i(z)H_{\text{mf}}(z)$
- GP *regression* then involves training these samples based on how well they fit the data

$$P(H(z) | D) = \int d\sigma_f d\ell d\phi \mathcal{L}(D | H(z)(\sigma_f, \ell, \phi)) P(\sigma_f, \ell, \phi) / P(D)$$


Testing Λ CDM via GP Hyperparameters

- These sort of tests can be performed because the hyperparameters of the GP regression encode information about whether the mean function is a good fit to the data
- i.e. how much information beyond the mean function is required to fit the data
- This test is performed by calculating the posterior of the hyperparameters to see if σ_f , the parameter that describes the heights of the fluctuations in the GP, is consistent with 0 or not
- If $\sigma_f > 0$ then, data need more flexibility than the given mean function
- If mean function standard model, the GP can test if the standard model is sufficient

Shafieloo A., Kim A. G., Linder E. V., 2012, Phys. Rev. D
Keeley, Shafieloo, L'Huillier, Linder MNRAS (2020)

Consistency with Λ CDM

- Hyperparameters of GP reconstruction
- Data : Pantheon SN and SDSS BAO
- Mean function : best fit LCDM to both data sets
- $\sigma_f = 0$ according to the data
- Therefore LCDM is consistent with the joint datasets




Keeley, Shafieloo, Zhao et al AJ 2021, arxiv:2010.03234

Model independent reconstruction of expansion history


$$h(z) = H(z)/H_0$$

- Orange - 68% and 95% CLs for LCDM
- Blue - Bands - 68% and 95% CLs; Lines - example GP reconstruction samples that have a better χ^2 than the best-fit LCDM



Keeley, Shafieloo, Zhao et al AJ 2021, arxiv:2010.03234


Tension Triangles



Keeley et al (in prep.)

Correlations

- Data: CMB+BAO+SN
- Colorbar: H_0
- Correlation between H_0 and Ω_m is simple primarily from $\Omega_m h^2$
- Correlation between w_0 and H_0 comes from θ_s



Keeley et al (in prep.)

Conclusions

- SN+BAO alone have **large degeneracies** in extended parameter spaces
- CMB alone has **large degeneracies** in extended parameter spaces
- BAO and SN are completely consistent with each other (in extended parameter spaces) and with LCDM
- BAO+SN are broadly consistent with Planck, even in extended parameter spaces
- **No preference** for curvature / evolving DE, but large swaths of parameter space are **still allowed**
- BAO+SN+CMB are in stark tension with SH0ES H_0 , even in extended parameter spaces
- Taken all together, new **low-redshift physics cannot solve the H_0 tension**