

Examination of Schrodinger equation in pre Planckian space-time early universe

Andrew Walcott Beckwith

Physics Department, Chongqing University, College of Physics, Chongqing University Huxi Campus, No. 44 Daxuechen Nanlu, Shapinba District, Chongqing 401331, People's Republic of China

Rwill9955b@gmail.com; abeckwith@uh.edu

Abstract

We look at Viutilli (1999) write up of a generalized schrodinger equation with its Ricci scalar inclusion, in curved space-time. This has a simplified version in Pre Planckian regime, which leads to comparing a resultant admissible wave function with Bohmian reformulations of quantum physics. As was done earlier, we compare this result with a formulation of a modified 'Poisson' equation from Poissons and Will from 2014, and then use inflaton physics . The resuting inflaton is then compared to the wavefunctional in the first part of this document.

Key words, Ricci tensor, Schrodinger equation, Modified Poisson Equation , massive gravity, inflaton physics.

1. Viutilli treatment of Schrodinger equation for curved space-time

Here, we bring up [1], and a re set of the Schrodinger equation in early space-time, with curvature.

In [1] we start with

$$i\hbar\partial_t\Psi = -\frac{\hbar^2}{2m}\nabla^2\Psi + mc^2\Psi \quad (1)$$

Through the following substitutions,

$$\nabla^2 \rightarrow \left(\frac{p^\alpha p^\beta}{m^2 c^2} - g_{\alpha\beta} \right) \nabla_\alpha \nabla_\beta$$

&

$$p^\alpha = i\hbar \cdot g^{\alpha\beta} \nabla_\beta$$

$$p_\beta = i\hbar \cdot \nabla_\beta$$

&

$$mc^2 = \frac{g^{\alpha\beta} p_\alpha p_\beta}{m} \quad (2)$$

Then, after more derivation [1] obtained, with using the Ricci tensor

R^c_a the following result:[2]

$$\nabla^a \nabla_a \left(\nabla^b \nabla_b + \frac{m^2 c^2}{\hbar^2} \right) \Psi + \nabla^a R^c_a \nabla_c \Psi = 0 \quad (3)$$

2. Simplifying Eq. (3) in Pre Planckian space-time

We will re write Eq. (3) to read as follows, with the result that in Pre Planckian-space-time.

$$\left(\partial_r^4 + \frac{m^2 c^2}{\hbar^2} \partial_r^2 \right) \Psi + R_r^r \partial_r^2 \Psi = 0 \quad (4)$$

Ricci tensor R^c_a in this setting becomes a constant, and is part of how the wave function evolves, and our candidate wave functional takes the form of

$$\Psi \propto \Psi_{initial} e^{sr} \sim \Psi_{initial} \exp \left(i \sqrt{\frac{m^2 c^2}{\hbar^2} + R_r^r} \cdot r \right) \quad (5)$$

We will be identifying what to put into [3] , i.e., the Ricci scalar as given in Pre Planckian space-time

$$\begin{aligned}
R_a^c \xrightarrow{\text{Pre-Planckian}} R_r^r &= 6 \cdot \left(\left[\frac{\ddot{a}}{a} \right] + \left[\frac{\dot{a}}{a} \right]^2 + \frac{k(\text{flatness})}{a^2} \right) \\
&= 12 \cdot H_{\text{initial}}^2 + \frac{6k(\text{flatness})}{a_{\text{initial}}^2} \sim 12 \cdot H_{\text{initial}}^2
\end{aligned} \tag{6}$$

We are assuming that the term $k(\text{flatness})$ approaches zero in Pre Planckian space-time. Obviously if it did not, the last term

$$\frac{6k(\text{flatness})}{a_{\text{initial}}^2} \quad \text{would be dominant.}$$

Having said this, with an evolutionary equation statement as to the phase value of Ψ , it is time to look at the initial Ψ_{initial} and to try to learn some physics from it.

3. Defining the initial value Ψ_{initial} of Eq. (5)

In order to do this, it may be useful to look at the classical degeneracy argument for forming Ψ_{initial} , and the reference by Rubakov may be useful for this [4] i.e. using a false vacuum analogy, we can write, if q is a generalized space-time unit of “length”, and we examine a quartic potential, i.e. look at

$$V(q) = -\frac{\mu^2 \cdot q^2}{2} + \frac{\tilde{\lambda} \cdot q^4}{4}$$

&

$$S(\text{action}) = \int_{q_0^-}^{q_0^+} \sqrt{2MV(q)} dq$$

$$\Rightarrow \Psi_{\text{initial}} \propto \Psi_0^+ \sim \exp\left(-\int_0^{q^+} dq \sqrt{2MV(q)}\right) \quad (7)$$

Now, use the usual given

$$\frac{M}{2} \frac{dq}{d\tau} = \frac{Mc}{2} = V(q)$$

$$\Rightarrow \Psi_{\text{initial}} \propto \Psi_0^+ \sim \exp\left(-\int_0^{q^+} dq M \sqrt{c}\right) \quad (8)$$

$$\xrightarrow{c=1} \Psi_0^+ \sim \exp(-Mq^+)$$

If using Ng infinite quantum statistics, [5], and a non zero massive graviton mass (massive gravity) [6]

$$E = M = [S(\text{entropy}) \sim n(\text{count} - \text{gravitons})] \cdot m_{\text{graviton}} \quad (9)$$

And if we have here that q^+ is in some sense proportional to length less than or equal to Planck length, the astonishing conclusion

is that Eq. (8) would probably be biased toward a low (nonzero) entropy count, which would mean for finite initial entropy, connected with information transfer, that we would through Eq. (8) and Eq. (9) have a bias toward initially low, say 10^5 or so initial entropy, as a way to quantify the input into the formation of an initial wavefunction, using entropy as equivalent to information as given by Lloyd[7]

4. Comparing the inputs into Eq(5), Eq.(6), Eq(8) and Eq. (9) against force against individual ‘gravitons’ in the Pre Planckian space-time

We will go to the [3] reference, page 85, in order to look at a change in the stress energy tensor,

$$\Delta T^{00} = (-T_{,j}^{0j} + F^{(0)}) \cdot \Delta t$$

$$F^{(0)} = \textit{force} / \textit{fluid}(\textit{unit} - \textit{volume}) \quad (10)$$

Using this, and stating that in the Pre Planckian regime of space-time due to [8], that $T_{,j}^{0j} \approx 0$, then if so, using [3] and [8]

$$F^{(0)} = \text{force} / \text{fluid}(\text{unit} - \text{volume})$$

$$T_{,j}^{0j} \approx 0$$

$$\Rightarrow \Delta T^{00} = (F^{(0)}) \cdot \Delta t$$

$$\Delta E = (F^{(0)}) \cdot \Delta V^{(4)} = \frac{1}{\Delta t \cdot a_{initial}^2 \phi}$$

$$\Rightarrow F^{(0)} = \frac{1}{\Delta V^{(3)} (\Delta t)^2 \cdot a_{initial}^2 \phi} \quad (11)$$

$$= \text{force} / \text{fluid}(\text{unit} - \text{volume})$$

Here, we have that the initial volume $\Delta V^{(3)}$ would be less than the cube power of a Planck length [9], but not zero, whereas the initial time would be less than, Planck time, but not equal to zero, [9]. For our analysis in the Pre Planckian regime, we will specify

$a_{initial}^2$ as the square of a nonzero initial scale factor for a nonsingular regime of space-time for General relativity with a value as given by [10], and elaborated upon in [11]. In addition we have that the inflaton, as given in Eq. (11) is explained in context by [12], and we will use an argument below as to how a nonzero graviton mass is linkable to a non zero initial radii, which in turn will state that

it is highly unlikely that $\Delta V^{(3)}$ which is proportional to a radial distance, cubed, goes to zero.. Note that Eq. (11) is also an argument as to why there would be a finite, not almost “infinite” initial value of entropy, so then that Eq. (8) would not go to zero.

Having said that, let us use a semi classical argument as to why the radii would not go to zero, even in Pre Planckian space-time. This

would be to insure that $\Delta V^{(3)}$ would not go to zero, even in Pre Planckian space-time

5. What is important about the modified Poissons equation [12]? Getting a non zero initial $\Delta V^{(3)}$

We will first of all refer to two necessary and sufficient conditions for the onset of a massive graviton given in [13] and combined with Padmanabhan's reference [12].

I.e., what we will be doing is to re do the reference calculations given in [13] with

$$\left(\nabla^2 + \left[\lambda^{-2} = \left(\frac{m_{\text{graviton}} c}{\hbar} \right)^2 \right] \right) \left[U = \frac{Gm}{r} \cdot \exp \left[(-r / \lambda) = \left(\frac{r \cdot m_{\text{graviton}} c}{\hbar} \right) \right] \right] = -4\pi G \rho$$

(12)

Here, we will be using in the Pre Planckian potential the inputs from the data usually associated with [12]

$$a \approx a_{\min} t^\gamma$$

$$\Leftrightarrow \phi \approx \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right\}$$

$$\Leftrightarrow V \approx V_0 \cdot \exp \left\{ -\sqrt{\frac{16\pi G}{\gamma}} \cdot \phi(t) \right\} \tag{13}$$

In other words, we will be using the inflation given by

$$\phi \approx \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right\} \quad (14)$$

If so, then

$$\left[\frac{d^2}{dr^2} + \left(\frac{m_{\text{graviton}} c}{\hbar} \right)^2 \right] \cdot \left(\frac{r^{-1} \alpha \cdot (3\alpha - 1)}{32\pi^2} \right) = G \cdot \rho \quad (15)$$

Then, after algebra, we have the following , from [14]

$$\left(m_{\text{graviton}} \right)^2 \approx \left[\frac{32\pi^2 r \cdot \hbar^2 \cdot G \cdot \rho}{c^2 \cdot \alpha \cdot (3\alpha - 1)} - \frac{16\pi^2 r^{-1} \cdot \hbar^2 \cdot G^{-2}}{c^2 \cdot \alpha \cdot (3\alpha - 1)} \right] \quad (16)$$

The quadratic Equation this engenders is, how to say.

$$\begin{aligned} \left(m_{\text{graviton}} \right)^2 &\approx \left[\frac{32\pi^2 r \cdot \hbar^2 \cdot G \cdot \rho}{c^2 \cdot \alpha \cdot (3\alpha - 1)} - \frac{16\pi^2 r^{-1} \cdot \hbar^2 \cdot G}{c^2 \cdot \alpha \cdot (3\alpha - 1)} \right] \\ \Rightarrow r^2 - \frac{r \cdot \left(m_{\text{graviton}} \right)^2}{\left(\frac{32\pi^2 \hbar^2 \cdot G \cdot \rho}{c^2 \cdot \alpha \cdot (3\alpha - 1)} \right)} - \frac{\left(\frac{16\pi^2 \cdot \hbar^2 \cdot G}{c^2 \cdot \alpha \cdot (3\alpha - 1)} \right)}{\left(\frac{32\pi^2 \hbar^2 \cdot G \cdot \rho}{c^2 \cdot \alpha \cdot (3\alpha - 1)} \right)} & \\ \doteq r^2 - \frac{r \cdot \left(m_{\text{graviton}} \right)^2}{\left(\frac{32\pi^2 \hbar^2 \cdot G \cdot \rho}{c^2 \cdot \alpha \cdot (3\alpha - 1)} \right)} - \frac{1}{2\rho} &= 0 \end{aligned} \quad (17)$$

A candidate for the density functional will come next, with the way of obtaining a critical value for r is given by [14] as follows, i.e. if

$$V_{Pre-Planckian} \sim \left(\Delta E \sim \frac{\hbar}{\delta t \cdot a_{\min}^2 \phi_{\inf}} \right) \quad (18)$$

As far as applications to:[1]

$$r^2 - \frac{r^4 \cdot (m_{graviton})^2 a_{\min}^2 \cdot \delta t \cdot \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right\}}{\left(\frac{32\pi^2 \hbar^2 \cdot G}{c^2 \cdot \alpha \cdot (3\alpha - 1)} \right)} = 0 \quad (19)$$

This would, lend itself to a quadratic equation for r, and the cube of r would be proportional to $\Delta V^{(3)}$ which would be non-zero, in Pre Planckian-space-time conditions.

6. Comparing the results of a non zero $\Delta V^{(3)}$ in Pre Planckian wave functional for Eq. (8) against De Broglie- Bohmian path of a wave functional

What we are examining if our qualitative argument which in sum yields a Pre Planckian wavefunction as

compared against the construction given in [15] which is in spirit comparable, up to a point with [16]

According to [15] we would have

$$\Psi(x,t) = \Psi(x_0,t_0) \exp \left[\left(\frac{i}{\hbar} \int_{x_0,t_0}^{x,t} dt \left[\frac{(\nabla S(action))^2}{2m} - (Q+V) \right] \right) - \int_{x_0,t_0}^{x,t} dt \frac{(\nabla^2 S(action))}{2m} \right] \quad (20)$$

Here, $S(action)$ would be the same as Eq. (7), whereas we have V as given by Eq. (18), and then we have

$$\begin{aligned} R &= \text{Re } \Psi(x,t) \\ Q &= \frac{-\hbar^2 \nabla^2 R}{2mR} \\ \nabla &\xrightarrow{\text{Pre-Planckian}} \partial_r \quad (21) \end{aligned}$$

Here, if we use $\hbar = c = 1$, and have $\Psi(x_0,t_0)$ as given by Eq. (8) whereas we find

$$\hbar = c = 1$$

$$Q = \frac{-\hbar^2 \nabla^2 R}{2mR} \xrightarrow{\text{Pre-Planckian}} \frac{1}{2M} \cdot (m_g^2 + R_r^r) \quad (22)$$

Here, M would be given by Eq. (9), i.e. and V were given by Eq. (18) we would find to a point that

$$\Psi(x,t) \sim \Psi(x_0,t_0) \exp \left[\left(\frac{i}{\hbar} \int_{x_0,t_0}^{x,t} dt \left[\frac{(\nabla S(action))^2}{2m} - (Q+V) \right] \right) \right] \quad (23)$$

I.e. the term $\Psi(x_0, t_0)$ would likely be the same, but interesting enough, $\int_{x_0, t_0}^{x, t} dt \frac{(\nabla^2 S(action))}{2m}$ would likely be almost zero, i.e. not contributing at all. i.e.

7. Conclusion, Overlap with the Bohmian quantum picture of physics if Eq. (24) is confirmed for Pre Planckian Space-time

The question to ask, is the following true ? This has to be confirmed rigorously.

$$\int_{x_0, t_0}^{x, t} dt \frac{(\nabla^2 S(action))}{2m} \xrightarrow{\text{Pre-Planckian}} \epsilon^+ \sim 0 \quad (24)$$

The significance of proving or falsifying Eq. (24) will in the end be part of future data analysis which should not contravene [17, 18, 19]. I.e. the experimentally implied limits should be adhered to and studied rigorously.

Furthermore, we have that analysis of Eq. (24) may be in tandem with analysis of the Corda paper [20] as to if gravity is possibly scalar-tensor, an extension of GR (possibly with some semi classical treatment of presumed quantum gravity formulations), or something else

Finally, since our paper is with respect to relic conditions, if so, then if we are to use a variant of interferometer methods, reference [21] and [22] if relic conditions are observable, via some form of space bound system, may allow us to with refinements get enough control of stochastic noise contamination of GW and the foot print of massive gravity to come up with confirmable data sets as to early universe

conditions. With luck, with considerable refinement of instrumentation, we may also be able to get experimental confirmation of [23] and its predictions as to inflaton physics and possibly massive gravity

6.. Acknowledgements

This work is supported in part by National Nature Science Foundation of China grant No. 11375279

References

- [1] Vuille, C. “Schrodingers’ Equation in General Relativity”, pp 243-246, in “General Relativity and Relativistic Astrophysics, Eighth Canadian Conference, Ed by Burgess, and Myers, R, AIP conference proceedings, 493, Melville, New York, USA 1999
- [2] Carmelli, M, “Classical Fields, General Relativity and Gauge Theory”:, Wiley Interscience, New York, New York, USA, 1982
- [3] Scott, R, “ A Student manual for A first course in General relativity”, Cambridge University Press, Cambridge UK, 2016
- [4] Rubakov, V, “ Classical theory of Gauge Fields”, Princeton University Press, Princeton, New Jersey, USA, 2002
- [5] Ng, Y.J. (2008) Spacetime Foam: From Entropy and Holography to Infinite Statistics and Nonlocality. Entropy, 10, 441-461. <http://dx.doi.org/10.3390/e10040441>
- [6] Goldhaber, A. and Nieto, M. (2010) Photon and Graviton Mass Limits. Reviews of Modern Physics, 82, 939-979. <http://arxiv.org/abs/0809.1003><http://dx.doi.org/10.1103/RevModPhys.82.939>

[7] Lloyd, S, "A theory of quantum gravity based on quantum computation", <https://arxiv.org/abs/quant-ph/0501135>

[8] Beckwith, A. (2016) Gedanken Experiment for Refining the Unruh Metric Tensor Uncertainty Principle via Schwarzschild Geometry and Planckian Space-Time with Initial Nonzero Entropy and Applying the Riemannian-Penrose Inequality and Initial Kinetic Energy for a Lower Bound to Graviton Mass (Massive Gravity). *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 106-124.
doi: [10.4236/jhepgc.2016.21012](https://doi.org/10.4236/jhepgc.2016.21012).

[9] Schwartz, B, "A first course in General Relativity, 2nd edition", Cambridge University Press, Cambridge, United Kingdom, 2011

[10] Camara, C.S., de Garcia Maia, M.R., Carvalho, J.C. and Lima, J.A.S. (2004) Nonsingular FRW Cosmology and Non Linear Dynamics. Arxiv astro-ph/0402311 Version 1, Feb 12.

[11] Beckwith, A. (2016) Thermodynamics and the Energy of Gravity Waves, GW's, as a Consequence of Non Linear Electrodynamics, NLED, Leading to an Initial Temperature and Strain h for Gravity Waves, GW, at the Start of Inflation. *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 98-105.
doi:[10.4236/jhepgc.2016.21011](https://doi.org/10.4236/jhepgc.2016.21011).

[12] Padmanabhan, T., "understanding our universe, current status and open issues", pp 175-204, of 100 Years of Relativity, Space-Time Structure: Einstein and Beyond, Editor, A. Ashatekar, World Scientific Publishing Co. Pte.Ltd, Singapore, Republic of Singapore, 2005; <http://arxiv.org/abs/gr-qc/0503107>

[13] Poisson, E., and Will, C., “Gravity, Newtonian , Post Newtonian, Relativistic”, Cambridge University Press, Cambridge, UK, 2014

[14] Beckwith, A. Accepted paper , for JHEPGC (approved for publication), “Examination of sufficient conditions for forming mass of “massive graviton”, from early universe”

[15] Licata, I, and Fiscaletti, D. “Quantum Potential: Physics, Geometry, and Algebra”, Springer-Verlag, New York, New York USA 2014

[16] Beckwith, A. (2016) Gedanken Experiment for Looking at δg_{tt} for Initial Expansion of the Universe and Influence on HUP via Dynamical Systems, with Positive Pre-Planckian Acceleration. *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 531-545. doi: [10.4236/jhepgc.2016.24046](https://doi.org/10.4236/jhepgc.2016.24046).

[17] [6] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration) ; “Observation of Gravitational Waves from a Binary Black Hole Merger”; Phys. Rev. Lett. **116**, 061102 – Published 11 February 2016

[18] [7] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration); “GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence”, Phys. Rev. Lett. **116**, 241103 – Published 15 June 2016

[19] [8] B. P. Abbott et al. “Tests of general relativity with GW150914”, <https://arxiv.org/pdf/1602.03841.pdf>

[20] Corda, C. “Interferometric detection of gravitational waves: the definitive test for General Relativity”, Int. J. Mod. Phys. D18:2275-2282,2009; <https://arxiv.org/abs/0905.2502>

[21] Willke, B, et al. ,“Stabilized High Power Laser for Advanced Gravitational Wave Detectors”,
<http://iopscience.iop.org/article/10.1088/1742-6596/32/1/040/meta;jsessionid=B072B9D72E7EB4F993D70061EEE365CF.c5.iopscience.cld.iop.org>

[22] Flaminico, R. et al. “A study of coating mechanical and optical losses in view of reducing mirror thermal noise in gravitational wave detectors” - .Class.Quant.Grav. 27 (2010) 084030

[23] Corda, C. “ Gravity’s Primordial Breath, ” Electronic Journal of Theoretical physics, EJTP 9, No. 26 (2012) 1–10; <http://www.ejtp.com/articles/ejtpv9i26p1.pdf>