## Sixteenth Marcel Grossmann Meeting



Contribution ID: 804

Type: Invited talk in the parallel session

## Lesson's from LIGO-Virgo's Biggest Black Holes

Thursday, 8 July 2021 16:50 (20 minutes)

Stellar theory predicts the existence of a black hole mass gap in the range ~50 to ~120 solar masses resulting from pair instability supernovae. The binary black holes of LIGO-Virgo's first two observing runs supported this prediction, showing evidence for a dearth of component black hole masses above 45 solar masses. Meanwhile, among the 30+ new observations from the third observing run, there are several black holes that appear to sit above the 45 solar mass limit. I will discuss how these unexpectedly massive black holes fit into our understanding of the binary black hole population. The data are consistent with several scenarios, including a mass distribution that evolves with redshift and the possibility that the most massive binary system, GW190521, straddles the mass gap, containing an intermediate-mass black hole heavier than 120 solar masses. I will also discuss applications of the binary black hole population to cosmology.

**Primary author:** FISHBACH, Maya (CIERA/ Northwestern University)

Presenter: FISHBACH, Maya (CIERA/ Northwestern University)Session Classification: Exploring the Black Hole Mass Gap

Track Classification: Black Holes: Theory and Observations/Experiments: Exploring the Black Hole

Mass Gap