



Contribution ID: 834

Type: Talk in the parallel session

Stress-energy Tensor for a Quantized Scalar Field When a Black Hole in Four Dimensions Forms From the Collapse of a Null Shell

Monday, 5 July 2021 17:55 (25 minutes)

A method will be presented which allows for the numerical computation of the stress-energy tensor for a quantized massless minimally coupled scalar field in the region outside the event horizon of a 4D Schwarzschild black hole that forms from the collapse of a null shell. This method involves taking the difference between the stress-energy tensor for the *in* state in the collapsing null shell spacetime and that for the Unruh state in Schwarzschild spacetime. The construction of the modes for the *in* vacuum state and the Unruh state will be discussed. Applying the method, the renormalized stress-energy tensor in the 2D case has been computed numerically and shown to be in agreement with the known analytic solution. In 4D, the presence of an effective potential in the mode equation causes scattering effects that make the construction of the *in* modes more complicated. The numerical computation of the *in* modes in this case will be presented.

Primary author: GHOLIZADEH SIAHMAZGI, Shohreh (Wake Forest University)

Co-authors: Dr ANDERSON, Paul R. (Wake Forest University); Mr CLARK, Raymond D. (Wake Forest University); Dr FABBRI, Alessandro (Universidad de Valencia-CSIC and Universit{e} Paris-Saclay)

Presenter: GHOLIZADEH SIAHMAZGI, Shohreh (Wake Forest University)

Session Classification: Black Hole Thermodynamics

Track Classification: Black Holes: Theory and Observations/Experiments: Black hole thermodynamics