## Sixteenth Marcel Grossmann Meeting



Contribution ID: 1019

Type: Talk in the parallel session

## Bardeen Black Hole from a Self-Dual Radius in Spacetime

Tuesday, 6 July 2021 11:00 (20 minutes)

From a phenomenological point of view, the singularities of ordinary black hole solutions hint at a breakdown of general relativity. The Bardeen black hole is a prototype of regular black hole solutions, i.e. those which are geodesically complete and avoid a curvature singularity.

The Bardeen solution is generally interpreted as a magnetically or electrically charged solution to gravity coupled to non-linear electrodynamics. In this talk, we derive that in a spacetime inheriting a self-dual radius from string theory, black holes naturally take on the Bardeen form.

The threshold mass for Bardeen black holes achieves a particular interpretation. During the evaporation process, a black hole undergoes a transition from an instable classical Schwarzschild phase to a stable quantum phase. In the end, there is a cold, thermodynamically stable remnant. The minimal black hole size of the order of the self-dual radius.

The self-dual radius can be tested by its impact on quantum mechanical systems. We find that it modifies the form of the atomic electrostatic potentials. We derive experimental bounds from high-precision spectroscopy of the hydrogen atom. The investigation of the  $1S_{1/2}-2S_{1/2}$  transition frequency allows to constrain the self-dual radius down to below  $3.9 \times 10^{-19}$  m.

**Primary author:** WONDRAK, Michael Florian (Goethe-Universität Frankfurt and Helmholtz Research Academy Hesse for FAIR )

**Presenter:** WONDRAK, Michael Florian (Goethe-Universität Frankfurt and Helmholtz Research Academy Hesse for FAIR )

Session Classification: Black Holes in Alternative Theories of Gravity

**Track Classification:** Black Holes: Theory and Observations/Experiments: Black Holes in alternative theories of gravity