



Contribution ID: 1009

Type: **Invited talk in the parallel session**

## **Tadpole contribution to magnetic photon-graviton conversion**

*Tuesday, 6 July 2021 09:50 (20 minutes)*

Photon-graviton conversion in a magnetic field is a process that is usually studied at tree level, but the one-loop corrections due to scalars and spinors have also been calculated. Differently from the tree-level process, at one-loop one finds the amplitude to depend on the photon polarization, leading to dichroism. However, previous calculations overlooked a tadpole contribution of the type that was considered to be vanishing in QED for decades but erroneously so as shown by Gies and Karbstein in 2016. Here we compute this missing contribution in closed form and study its numerical relevance compared to the standard one.

**Primary authors:** SCHUBERT, Christian (Instituto de Fisica y Matematicas, Universidad Michoacana de San Nicolas de Hidalgo); Dr AHMADINIAZ, Naser (Helmholtz-Zentrum Dresden-Rossendorf); Prof. BASTIANELLI, Fiorenzo (Dipartimento di Fisica ed Astronomia, Universita di Bologna, and INFN, Sezione di Bologna); KARBSTEIN, Felix

**Presenters:** SCHUBERT, Christian (Instituto de Fisica y Matematicas, Universidad Michoacana de San Nicolas de Hidalgo); KARBSTEIN, Felix

**Session Classification:** Strong Electromagnetic and Gravitational Field Physics: From Laboratories to Early Universe

**Track Classification:** Strong Field: Strong Electromagnetic and Gravitational Field Physics: From Laboratories to Early Universe