On quasinormal modes in 4D black hole solutions in the model with anisotropic fluid

Thursday, 15 June 2023 17:30 (15 minutes)

We consider a family of 4-dimensional black hole solutions

governed by natural number q = 1, 2, 3, ..., which appear in the model with anisotropic fluid and the equations of state: $p_r = -\rho(2q - 1)^{-1}$, $p_t = -p_r$, where p_r and p_t are pressures in radial and transverse directions, respectively, and $\rho > 0$ is the density. These equations of state obey weak, strong and dominant energy conditions. For q = 1 the metric of the solution coincides with that of the Reissner-Nordstr\"om one. The global structure of solutions is outlined, giving rise to Carter-Penrose diagram of Reissner-Nordstr\"om or Schwarzschild types for odd q = 2k + 1 or even q = 2k, respectively. Certain physical parameters corresponding to BH solutions

(gravitational mass, PPN parameters, Hawking temperature and entropy) are calculated. We obtain and analyse the quasinormal modes for a test massless scalar field in the eikonal approximation. For limiting case $q = +\infty$, they coincide with the well-known results for the Schwarzschild solution. We show that the Hod conjecture which connect the Hawking temperature and the damping rate is obeyed for all $q \ge 2$ and all (allowed) values of parameters.

Primary authors: Dr BOLOKHOV, Sergey (RUDN); IVASHCHUK, Vladimir (RUDN/VNIIMS)

Presenter: IVASHCHUK, Vladimir (RUDN/VNIIMS)

Session Classification: Thursday afternoon session