Contribution ID: 13

Probing electromagnetic-gravitational wave emission coincidence in type I binary-driven hypernova family of long GRBs at very-high redshift (online talk)

Monday, 19 June 2023 11:00 (45 minutes)

Due to the technical time delay of the XRT instrument on board the Neil Gehrels Swift Observatory satellite, we cannot observe the X-ray emission occurring less than ~40 s after a gamma-ray burst (GRB) trigger time. We here indicate a new strategy of using the cosmological time dilatation in high redshift GRBs to observe the earliest X-ray emission by Swift/XRT in the GRB cosmological rest-frame. We illustrate this procedure using 354 GRBs with a well-defined cosmological redshift selected from the Swift GRB catalog. We compare and contrast the time delay between the trigger of the source and the first observation by Swift/XRT as measured in the observer frame (OTD) and the corresponding delay measured in GRBs' cosmological rest-frame (RTD). We consider as specific prototypes three binary-driven hypernovae of type I (BdHNe I): GRB 090423 at z=8.2 with an RTD of 8.2 s, GRB 090429B at z~9.4 with an RTD of 10.1 s, as well as the GRB 220101A at z=4.6 with an RTD of 14.2 s. This opens a new possibility for probing Episode (1) of BdHNe, linked to the origin and early appearance of the newborn neutron star (vNS) and its transition from a Jacobi triaxial ellipsoid (JTE) to a Maclaurin spheroid configuration that originates the GRB afterglow onset. We also present the methodology to compute the sweeping frequencies and the energetics of the associated conspicuous gravitational wave emission.

Primary authors: BIANCO, Carlo Luciano (ICRANet); RASTEGARNIA, F.; RUEDA, J.A.; LI, L.; DELLA VALLE, M.; MIRTORABI, M.T.; MORADI, R.; RUFFINI, R.; ZHANG, S.R.; WANG, Y.

Presenter: BIANCO, Carlo Luciano (ICRANet)