



Contribution ID: 429

Type: **Talk in a parallel session**

## **Non-thermal radiation due to CR transport in the magnetic halo of the Milky Way**

*Tuesday, 9 July 2024 18:00 (20 minutes)*

The large-scale structures such as Fermi Bubbles and eROSITA Bubbles provide a unique opportunity to study our Milky Way. However, the nature and origin of these large structures are still under debate. In this talk, I will present the identification of several kpc-scale magnetised structures based on their polarized radio emission and their gamma-ray counterparts, which can be interpreted as the radiation of relativistic electrons in the Galactic magnetic halo. These non-thermal structures extend far above and below the Galactic plane and are spatially coincident with the thermal X-ray emission from the eROSITA Bubbles. Multi-wavelength spectral energy distribution analyses have revealed that both thermal and non-thermal extended structures can be explained by Galactic outflows driven by the active star-forming regions located at 3 – 5 kpc from the Galactic Centre. These results reveal how X-ray-emitting and magnetised halos of spiral galaxies can be related to intense star formation activities.

**Primary author:** ZHANG, Heshou (INAF - OA Brera)

**Co-authors:** Dr PONTI, Gabriele (INAF - OA Brera); Dr CARRETTI, Ettore (INAF - IRA); Prof. LIU, Ruo-Yu (Nanjing University); Prof. MORRIS, Mark (UCLA); Prof. HAVERKORN, Marijke (Radboud University Nijmegen); Dr LOCATELLI, Nicola (INAF - OA Brera); Ms ZHENG, Xueying (MPE); Prof. AHARONIAN, Felix (Dublin Institute for Advanced Studies); Dr ZHANG, Hai-Ming (Nanjing University); Ms ZHANG, Yi (MPE); Mr STEL, Giovanni (INAF - OA Brera); Prof. STRONG, Andrew (MPE); Mr YEUNG, Micheal (MPE); Dr MERLONI, Andrea (MPE)

**Presenter:** ZHANG, Heshou (INAF - OA Brera)

**Session Classification:** High energy astrophysics

**Track Classification:** High energy (HE): High energy astrophysics