



Contribution ID: 222

Type: **Invited talk in a parallel session**

## A Cosmological Fireball with Sixteen-Percent Gamma-Ray Radiative Efficiency

*Monday, 8 July 2024 17:15 (15 minutes)*

Gamma-ray bursts (GRBs) are the most powerful explosions in the universe. How efficiently the jet converts its energy to radiation is a long-standing problem and it is poorly constrained. The standard model invokes a relativistic fireball with a bright photosphere emission component. A definitive diagnosis of GRB radiation components and measurement of GRB radiative efficiency require prompt emission and afterglow data with high-resolution and wide-band coverage in time and energy. Here we report a comprehensive temporal and spectral analysis of the TeV-emitting bright GRB 190114C. Its fluence is one of the highest of all GRBs detected so far, which allows us to perform a high-resolution study of the prompt emission spectral properties and their temporal evolution down to a timescale of about 0.1 s. We observe that each of the initial pulses has a thermal component contributing  $\sim 20\%$  of the total energy, the corresponding temperature and the inferred Lorentz factor of the photosphere evolve following broken power-law shapes. From the observation of the non-thermal spectra and the light-curve, the onset of afterglow corresponding to the deceleration of the fireball is considered at  $\sim 6$ -s. By incorporating the thermal and the non-thermal observations, as well as the photosphere and the synchrotron radiative mechanisms, we can directly derive the fireball energy budget with little dependence on hypothetical parameters and to measure a  $\sim 16\%$  radiative efficiency for this GRB. With the fireball energy budget derived, the afterglow microphysics parameters can also be constrained directly from the data.

**Primary authors:** Mr TIRADO, Alberto J. Castro (Instituto de Astrofísica de Andalucía (IAA-CSIC), PO Box 03004, 18008 Granada, Spain); Mr PEER, Asaf (Department of Physics, Bar-Ilan University, Ramat-Gan 52900, Israel); Mr ZHANG, Bing (Department of Physics and Astronomy, University of Nevada, Las Vegas, NV 89154, USA); Mr RYDE, Felix (Department of Physics, KTH Royal Institute of Technology, and the Oskar Klein Centre for Cosmoparticle Physics, 10691 Stockholm, Sweden); Ms PAGE, Kim (School of Physics and Astronomy, University of Leicester, University Road, Leicester LE1 7RH, UK); LI, Liang (ICRANet, Piazza della Repubblica 10, I-65122 Pescara, Italy); Mr BHAT, P.N. (Center for Space Plasma and Aeronomic Research, University of Alabama in Huntsville, Huntsville, AL, USA); Mr VERES, Peter (Center for Space Plasma and Aeronomic Research, University of Alabama in Huntsville, Huntsville, AL, USA); Mr GUIRIEC, Sylvain; Mr WU, Yang (ICRANet, Piazza della Repubblica 10, 65122 Pescara, Italy)

**Presenter:** LI, Liang (ICRANet, Piazza della Repubblica 10, I-65122 Pescara, Italy)

**Session Classification:** Emission mechanisms in gamma-ray bursts

**Track Classification:** Gamma-Ray Bursts (GB): Emission mechanisms in gamma-ray bursts